




Fig 1




}





F14.3.




F16. 4.















FIG BA.







d fi

F.g. 13




Figure 14




Figure 15




Figure 16: Calibration curve for fluorescein using Card. Experimental conditions: buffer 10.0 mM CHES pH=10.0, and 1% PEO; voltage configurations as shown in Table; PMT voltage, 400 V;




Figure 17: Alkaline phosphatase assay using Card, Effect of the Incubation Time. Experimental conditions: buffer 10.0 mM CHES pH=10.0, and 1% PEO; voltage configurations as shown in Table; PMT voltage, 800 V; Alkaline phosphatase was 1.33 mU/ml; FDP concentration, 3.33 μ M.




Figure 18: Calibration curve for alkaline phosphatase enzyme. Experimental conditions: buffer 10.0 mM CHES pH=10.0, and 1% PEO; voltage configurations as shown in Table; PMT voltage, 500 V; FDP concentration, 3.33 μ M




Figure 19: Alkaline Phosphatase Assay using Card. Effect of the Inhibitor Concentration (PNPP). Experimental conditions: buffer 10.0 mM CHES pH=10.0, and 1% PEO; voltage configurations as shown in Table; PMT voltage, 800 V; Alkaline phosphatase was 1.33 mU/ml; FDP concentration, 3.33 μ M and different concentration of PNPP as shown in the Figure; incubation time, 10 min.




Figure 20: Alkaline Phosphatase assay using Card. Calibration Curve for PNPP. Experimental conditions: buffer 10.0 mM CHES pH=10.0, and 1% PEO; voltage configurations as shown in Table; PMT voltage, 800 V; Alkaline phosphatase concentration was 1.33 mU/ml; FDP concentration, 3.33 μ M.